

Anatomic Coracoclavicular Reconstruction Rehabilitation

PHASE I - IMMEDIATE POSTSURGICAL PHASE (DAYS 1 - 14)

Goals

- 1. Maintain integrity of the repair
- 2. Diminish pain and inflammation
- 3. Prevent muscular inhibition

Days 1 - 14

"Gunslinger" brace

- Pendulum exercises 4x per day
- Elbow/hand gripping and ROM exercises
- Cryotherapy/ice for pain and inflammation 15-20 minutes every hour
- Sleeping sleep in sling or brace

Precautions To Be Maintained Through Week 6:

- 1. Maintain arm in brace, remove only for exercise
- 2. No lifting of objects
- 3. No excessive or aggressive stretching or sudden movements
- 4. No supporting of body weight by hands
- 5. No bimanual activities (pulling on pants/socks, etc.)

PHASE II - PROTECTION PHASE (WEEKS 3 - 6)

Goals

- 1. Allow healing of soft tissue
- 2. Do not overstress healing tissue
- 3. Gradually restore full PROM by weeks 4 6
- 4. Decrease pain and inflammation

Weeks 3 - 6

- Continue pendulum exercises
- Start formal physical therapy (PT)
- ° AAROM, gentle PROM as tolerated to match uninvolved shoulder
- · Continue use of cryotherapy/ice as needed if still swollen

PHASE III - INTERMEDIATE PHASE (WEEKS 7 - 12)

Goals

- 1. Maintain full A/AA/PROM
- 2. Dynamic shoulder stability

Weeks 7 - 12

- Discontinue use of brace/sling
- Continue AA/PROM
- Work on scapulohumeral rhythm
- May use arm for light activities of daily living (ADLs)
- May begin gentle AROM
- · No lifting or strengthening



Pendulum, Circular-Bend forward 90 degrees at the waist, using a table for support.

Rock body in a circular pattern to move arm clockwise 10 times, then counterclockwise 10 times. Do 3 sessions a day.

*American Academy of Orthopaedic Surgeons

PHASE IV - EARLY STRENGTHENING PHASE (WEEKS 13 - 16)

Goals

- 1. Maintain full nonpainful ROM
- 2. Enhance functional use of upper extremity
- 3. Improve muscular strength and power

Weeks 13 - 16

- Continue ROM and stretching to maintain full ROM
- Self capsular stretches
- Isometric strengthening only

PHASE V - PROGRESSIVE STRENGTHENING (WEEKS 17 - 23)

Goals

- 1. Maintain full nonpainful ROM
- 2. Enhance functional use of upper extremity
- 3. Improve dynamic muscular strength

Week 17 - 23

- · Continue stretching if motion is tight
- Begin dynamic strengthening and scapular strengthening (no push-ups)
- May initiate interval sport program (i.e., golf, etc.)

PHASE VI - RETURN TO UNRESTRICTED ACTIVITY (WEEKS 24+)

Goals

- 1. Maintain full nonpainful ROM
- 2. Enhance functional use of upper extremity
- 3. Regain full strength and unrestricted activity

Weeks 24+

- Continue stretching if motion is tight
- Continue dynamic strengthening and scapular strengthening
- May initiate contact sports